

Effects of the interaction between dark energy and dark matter on cosmological parameters

Jian-Hua He, Bin Wang*

Department of Physics, Fudan University, Shanghai 200433, P. R. China

Abstract

We examine the effects of possible phenomenological interactions between dark energy and dark matter on cosmological parameters and their efficiency in solving the coincidence problem. We work with two simple parameterizations of the dynamical dark energy equation of state and the constant dark energy equation of state. Using observational data coming from the new 182 Gold type Ia supernova samples, the shift parameter of the Cosmic Microwave Background given by the three-year Wilkinson Microwave Anisotropy Probe observations, and the baryon acoustic oscillation measurement from the Sloan Digital Sky Survey, we perform a statistical joint analysis of different forms of phenomenological interaction between dark energy and dark matter.

PACS numbers: 98.80.Cq, 98.80-k

arXiv:0801.4233v2 [astro-ph] 23 May 2008

* E-mail address: wangb@fudan.edu.cn

I. INTRODUCTION

Our universe is believed undergoing an accelerated expansion driven by a yet unknown dark energy (DE) [1]-[5]. Much effort has been devoted to understand the nature and the origin of the DE. The leading interpretation of such a DE is a cosmological constant with equation of state (EoS) $w = -1$. Although this interpretation is consistent with observational data, at the fundamental level it fails to be convincing. The vacuum energy density falls far below the value predicted by any sensible quantum field theory, and it unavoidably leads to the coincidence problem, i.e., “why are the vacuum and matter energy densities of precisely the same order today?”. To overcome the coincidence problem, some sophisticated dynamical DE models either relating the DE to a scalar field called quintessence with $w > -1$, or to an exotic field called phantom with $w < -1$ [6] have been put forward to replace the cosmological constant. But it is doubtful that there is a clear winner in sight to explain the DE and solve the coincidence problem.

DE contributes a significant fraction of the content of the universe, it is natural to consider its interaction with the remaining fields of the Standard Model in the framework of field theory. The possibility that DE and DM can interact has been studied in [7]-[16], among others. It has been shown that the coupling between a DE (or quintessence) field and DM can provide a mechanism to alleviate the coincidence problem [7, 10]. A suitable choice of the coupling, motivated by holographic arguments, can also lead to the crossing of the phantom barrier which separates models with EoS $w > -1$ from models with $w < -1$ [11, 12]. In addition, it has been argued that an appropriate interaction between DE and DM can influence the perturbation dynamics and affect the lowest multipoles of the CMB spectrum [13, 17]. Arguments using structure formation of galaxies suggested that the strength of the coupling could be as large as the QED fine structure constant [13, 18]. More recently, it was shown that such an interaction could be inferred from the expansion history of the universe, as manifested in, e.g., the supernova data together with CMB and large-scale structure [19]. Nevertheless, the observational limits on the strength of such an interaction remain weak [20]. In addition, it was suggested that the dynamical equilibrium of collapsed structures would be affected by the coupling of DE to DM [21, 22]. The basic idea is that the virial theorem is distorted by the non-conservation of mass caused by the coupling. This problem has been analyzed precisely in [23].

The interaction between DE and DM could be a major issue to be confronted in studying the physics of DE. However, due to the nature of these two components remain unknown, it will not be possible to derive the precise form of the interaction from first principles. One has to assume a specific coupling from the outset [24, 25] or determine it from phenomenological requirements [26]. In view of the continuity equations, the interaction between DE and DM must be a function of the energy densities multiplied by a quantity with units of inverse of time. For the latter the obvious choice is the Hubble factor H . Thus, the interaction between DE and DM could be expressed phenomenologically in forms such as $Q = Q(H\rho_{DM})$ [20, 36], $Q = Q(H\rho_{DE})$ [28], $Q = Q(H(\rho_{DE} + \rho_{DM}))$ [9]-[13] or most generally in the form $Q = Q(H\rho_{DE}, H\rho_{DM})$ which leads $Q \simeq \delta_1 H\rho_{DE} + \delta_2 H\rho_{DM}$ from the first term in the power law expansion. Considering the couplings are terms in the lagragian which mix both DE and DM, one may further presume that they could be parameterized by some product of the densities of DE and DM, for example $Q = \lambda\rho_{DE}\rho_{DM}$ [14]. Besides these phenomenological descriptions of the interaction between DE and DM, recently there is an attempt to describe the coupling from the thermodynamical consideration [27, 28].

It is of great interest to investigate effects of the interaction between DE and DM on the universe evolution, especially the influence of different forms of the interaction on cosmological parameters. This is the main motivation of the present paper. Using the new 182 Gold type Ia supernova samples (SNIa), the baryon acoustic oscillation measurement from the Sloan Digital Sky Survey (SDSS-BAO) and the shift parameter of the Cosmic Microwave Background given by the three-year Wilkinson Microwave Anisotropy Probe observations (CMB shift), we are going to examine the efficiency of different coupling forms in solving the cosmic coincidence problem. In our study we will not specify any special model of DE. Considering recent accurate data analysis showing that the time varying DE gives a better fit than a cosmological constant and in particular, DE EoS can cross -1 around $z = 0.2$ from above to below [29], we will employ two commonly used parameterizations in our work, namely $\omega_I = \omega_0 + \omega_1 z / (1 + z) = \omega_0 + \omega_1 (1 - a)$ and $\omega_{II} = \omega_0 + \omega_1 z / (1 + z)^2 = \omega_0 + (a - a^2)\omega_1$. For comparison we will also examine the effects of the interaction between DE and DM for the constant EoS of DE.

II. PHENOMENOLOGICAL MODELS DESCRIBING THE INTERACTION BETWEEN DE AND DM

In the FRW universe, the Friedmann equation reads

$$3M_p^2 H^2 = \rho_r + \rho_b + \rho_{DM} + \rho_{DE}. \quad (1)$$

Using the critical energy density $\rho_c^0 = 3M_p^2 H_0^2$, we can rewrite the Friedmann equation in the form

$$E^2(z) = \frac{H^2(z)}{H_0^2} = \Omega_b^0(1+z)^3 + \Omega_r^0(1+z)^4 + \rho_{DE}(z)/\rho_c^0 + \rho_{DM}(z)/\rho_c^0. \quad (2)$$

The energy density of the radiation is given by,

$$\rho_r^0 = \sigma_b T_{cmb}^4 \quad (3)$$

where σ_b is the Stefan-Boltzmann constant, $T_{cmb} = 2.726K$ is the CMB temperature at present. $\rho_{DE}(z)$ represents the DE density and ρ_{DM} is the energy density of the cold DM.

When DE and DM interact with each other, we have

$$\dot{\rho}_{DM} + 3H\rho_{DM} = Q, \quad (4)$$

$$\dot{\rho}_{DE} + 3H(1+w_{DE})\rho_{DE} = -Q, \quad (5)$$

respectively, where Q denotes the interaction term. Notice that the overall energy density of dark sectors $\rho_{DM} + \rho_{DE}$ is conserved.

From the equations above, phenomenological forms of the interaction between DE and DM must be a function of the energy densities multiplied by a quantity with units of inverse of time. The obvious choice of the quantity with units of inverse of time is the Hubble factor H . The energy densities could be chosen as density of either DE, DM, DE plus DM, the linear combination of DE and DM or the product of DE and DM.

We will compare these different phenomenological expressions of the interaction between DE and DM and investigate their effects on cosmological evolutions and efficiencies in solving the coincidence problem.

A. The coupling is proportional to the energy density of DE

If the interaction term is proportional to the energy density of DE, we choose $Q = \delta H \rho_{DE}$ and rewrite the continuity equations 4 and 5

$$\dot{\rho}_{DM} + 3H\rho_{DM} - \delta H \rho_{DE} = 0 \quad (6)$$

$$\dot{\rho}_{DE} + 3H\rho_{DE}(1+\omega) + \delta H \rho_{DE} = 0 \quad (7)$$

Employing the first commonly used parameterization of the EoS ω_I , we obtain a closed form solution for (7)

$$\rho_{DE} = \rho_{DE}^0 e^{-3\omega_1} e^{3\omega_1/(1+z)} (1+z)^{3(1+\omega_0+\omega_1)+\delta}. \quad (8)$$

Insert this solution into Eq(6), we get

$$\rho_{DM} = (1+z)^3 \rho_{DM}^0 - \delta \rho_{DE}^0 (1+z)^3 \int_0^z e^{3\omega_1/(1+z)-3\omega_1} (1+z)^{3(\omega_0+\omega_1)+\delta-1} dz. \quad (9)$$

Choosing the second parameterization of the EoS ω_{II} , we have

$$\rho_{DE} = \rho_{DE}^0 e^{\frac{3}{2}\omega_1 + \frac{3}{2}\omega_1/(1+z)^2 - 3\omega_1/(1+z)} (1+z)^{3(1+\omega_0)+\delta}, \quad (10)$$

and

$$\rho_{DM} = \rho_{DM}^0 (1+z)^3 - \delta \rho_{DE}^0 (1+z)^3 e^{\frac{3}{2}\omega_1} \int_0^z e^{\frac{3}{2}\omega_1/(1+z)^2 - 3\omega_1/(1+z)} (1+z)^{\delta+3\omega_0-1} dz. \quad (11)$$

B. The coupling is proportional to the energy density of DM

Expressing the interaction between DE and DM in proportional to the DM energy density $Q = \delta H \rho_{DM}$, we can rewrite the continuity equations as

$$\dot{\rho}_{DM} + 3H\rho_{DM} - \delta H \rho_{DM} = 0, \quad (12)$$

$$\dot{\rho}_{DE} + 3H\rho_{DE}(1+\omega) + \delta H \rho_{DM} = 0. \quad (13)$$

Employing ω_I as the parameterization of the EoS, from Eq(12) we obtain,

$$\rho_{DM} = \rho_{DM}^0 (1+z)^{3-\delta}. \quad (14)$$

Insert this ρ_{DM} into Eq(13), we arrive at

$$\rho_{DE} = (1+z)^{3(1+\omega_0+\omega_1)} e^{3\omega_1/(1+z)} \delta \rho_{DM}^0 \int_0^z e^{-3\omega_1/(1+z)} (1+z)^{-3(\omega_0+\omega_1)-\delta-1} dz \quad (15)$$

$$+ \rho_{DE}^0 (1+z)^{3(1+\omega_0+\omega_1)} e^{3\omega_1/(1+z)-3\omega_1}. \quad (16)$$

Using ω_{II} as the EoS, we find the evolution equation for DM as

$$\rho_{DM} = \rho_{DM}^0 (1+z)^{3-\delta}. \quad (17)$$

Then from Eq(13) we have the evolution of the DE reads

$$\rho_{DE} = (1+z)^{3(1+\omega_0)} e^{-\frac{3\omega_1}{1+z} + \frac{3}{2} \frac{\omega_1}{(1+z)^2}} \delta \rho_{DM}^0 \int_0^z (1+z)^{-3\omega_0-\delta-1} e^{\frac{3\omega_1}{1+z} - \frac{3}{2} \frac{\omega_1}{(1+z)^2}} dz \quad (18)$$

$$+ (1+z)^{3(1+\omega_0)} e^{-\frac{3\omega_1}{1+z} + \frac{3}{2} \frac{\omega_1}{(1+z)^2}} e^{\frac{3}{2}\omega_1} \rho_{DE}^0. \quad (19)$$

C. The coupling is proportional to the total energy density of DE and DM

We can also write the interaction between DE and DM in proportion to the total energy densities of DE and DM, so that the continuity equations read

$$\dot{\rho}_{DM} + 3H\rho_{DM} - \delta H(\rho_{DM} + \rho_{DE}) = 0, \quad (20)$$

$$\dot{\rho}_{DE} + 3H\rho_{DE}(1+\omega) + \delta H(\rho_{DM} + \rho_{DE}) = 0. \quad (21)$$

The above equations can be converted to

$$\frac{d\rho_{DM}}{dz} - \frac{3}{1+z}\rho_{DM} + \frac{\delta}{1+z}(\rho_{DE} + \rho_{DM}) = 0, \quad (22)$$

$$\frac{d\rho_{DE}}{dz} - \frac{3}{1+z}(1+\omega)\rho_{DE} - \frac{\delta}{1+z}(\rho_{DE} + \rho_{DM}) = 0. \quad (23)$$

D. The coupling is proportional to the linear combination of energy densities of DE and DM

The phenomenological coupling between DE and DM could be expressed most generally in the form $Q = Q(H\rho_{DE}, H\rho_{DM})$ which leads $Q \simeq \delta_1 H\rho_{DE} + \delta_2 H\rho_{DM}$ from the first term in the power law expansion. Thus the continuity equations read

$$\dot{\rho}_{DM} + 3H\rho_{DM} - H(\delta_1\rho_{DE} + \delta_2\rho_{DM}) = 0, \quad (24)$$

$$\dot{\rho}_{DE} + 3H\rho_{DE}(1+\omega) + H(\delta_1\rho_{DE} + \delta_2\rho_{DM}) = 0, \quad (25)$$

which can be changed into

$$\frac{d\rho_{DM}}{dz} - \frac{3}{1+z}\rho_{DM} + \frac{1}{1+z}(\delta_1\rho_{DE} + \delta_2\rho_{DM}) = 0, \quad (26)$$

$$\frac{d\rho_{DE}}{dz} - \frac{3}{1+z}(1+\omega)\rho_{DE} - \frac{1}{1+z}(\delta_1\rho_{DE} + \delta_2\rho_{DM}) = 0. \quad (27)$$

For the last two cases of describing the interaction between DE and DM, it is not easy to write out analytic solutions of ρ_{DE} and ρ_{DM} when we substitute the parameterizations of EoS. We will completely count on the numerical calculation to investigate the time evolution of the DE and DM in these cases.

E. The coupling is proportional to the product of energy densities of DE and DM

The coupling can be considered in terms of the lagrangian which mix both DE and DM, we can adopt the form of the interaction as the product of the densities of DE and DM [14], $Q = \lambda \rho_{DE} \rho_{DM}$, where $\lambda = kM^{-3}$ and M is considered as a mass parameter and k is a dimensionless constant. Taking $\Omega_x = \rho_x/\rho_{c0}$ and $\rho_{c0} = 3H_0^2 M_p^2/(8\pi) = h^2 \cdot 0.81 \cdot 10^{-46} \text{GeV}^4$, the dimensionless continuity equations for the DM and DE read

$$\begin{aligned} \frac{d\Omega_{DM}}{dz} - \frac{3}{1+z}\Omega_{DM} + \delta \frac{1}{(1+z)} \frac{\Omega_{DM}\Omega_{DE}}{\sqrt{\Omega_{DM} + \Omega_{DE}}} &= 0 \\ \frac{d\Omega_{DE}}{dz} - \frac{3}{1+z}(1+\omega)\Omega_{DE} - \delta \frac{1}{(1+z)} \frac{\Omega_{DM}\Omega_{DE}}{\sqrt{\Omega_{DM} + \Omega_{DE}}} &= 0 \end{aligned} \quad (28)$$

where $\delta = \sqrt{3/(8\pi)}\lambda\rho_{c0}^{1/2}M_p = \sqrt{3/(8\pi)}k(\frac{M_p}{M})(\frac{\rho_{c0}}{M^4})^{1/2}$, which is a dimensionless coupling constant now.

So far we have listed all possible phenomenological descriptions of the interaction between DE and DM. In the next section, we will examine the viability of these descriptions by comparing with observations.

III. OBSERVATIONAL CONSTRAINTS

We will use the three-year WMAP (WMAP3) data[30], the SN Ia data [31] and the Baryon Acoustic Oscillation (BAO) measurement from the Sloan Digital Sky Survey [32] to study the property of the interaction between DE and DM. In [33], the authors showed that combining the shift parameters R and the angular scale l_a of the sound horizon at recombination appears to be a good approximation of the full WMAP3 data. The model independent constraints on R and l_a was later given [34] by using the WMAP3 data, where the covariance matrix of the parameters R, l_a and $\Omega_b h^2$ was also provided. In the following we will use the shift parameter R , the angular scale l_a of the sound horizon at recombination and their covariance matrix given in [34] and employ the Monte-Carlo Markov Chain (MCMC) method to explore the parameter space. Our MCMC code is based on the publicly available package CMBeasy[38].

For the SNIa data, we will calculate $\chi^2 = \sum_i \frac{[\mu_{obs}(z_i) - \mu(z_i)]^2}{\sigma_i^2}$, where the extinction-corrected distance modulus $\mu(z) = 5\log_{10}[d_L(z)/\text{Mpc}] + 25$, σ_i is the total uncertainty in the SN Ia data, and the luminosity distance is $d_L(z) = (1+z)/H_0 \int_0^z dz'/E(z')$ and the nuisance parameter H_0 is marginalized over with flat prior.

For the SDSS data, we add the BAO parameter [30, 32],

$$A = \frac{\sqrt{\Omega_m^0}}{E(0.35)^{1/3}} \left[\frac{1}{0.35} \int_0^{0.35} \frac{dz'}{E(z')} \right]^{2/3} = 0.469(0.95/0.98)^{-0.35} \pm 0.017,$$

to χ^2 .

For WMAP3 data, we first add the shift parameter[34]

$$R = \sqrt{\Omega_m^0} \int_0^{z_{ls}} \frac{dz'}{E(z')} = 1.70 \pm 0.03,$$

to χ^2 at $z = 1089 \pm 1$. It was argued in [33, 34] that the combination of the shift parameter and the angular scale of the sound horizon at recombination gives much better constraints on cosmological parameters. Thus we add the angular scale of the sound horizon at recombination

$$l_a = \frac{\pi R / \sqrt{\Omega_m^0}}{\int_{z_{ls}}^{\infty} dz c_s / E(z)} = 302.2 \pm 1.2,$$

where the sound speed $c_s = 1/\sqrt{3(1+R_b a)}$, $R_b = 315000\Omega_b h^2 (T_{cmb}/2.7K)^{-4}$, a is the scale factor, and $\Omega_b h^2 = 0.022^{+0.00082}_{-0.00082}$.

We use the MCMC method to explore the parameter space. We treat H_0 as also a fitting parameter, and we impose a prior of $H_0 = 72 \pm 8 \text{km/s/Mpc}$ [35]. By using two parameterizations for the EoS of DE ω_I, ω_{II} , we obtain the parameter space for different couplings respectively and we will show our results in the following sections. For comparisons we will also present the result for constant DE EoS.




Figure 1: The likelihood for coupling parameters when the interaction is chosen as $\delta H \rho_{DE}$

A. The coupling is proportional to the energy density of DE

Choosing the first parameterization of the DE EoS ω_I and expressing the coupling between DE and DM in proportional to the energy density of DE, the results are shown in Fig 1a and Fig 2a. The quantity of interest for analyzing the coincidence problem is the ratio $r = \rho_{DM}/\rho_{DE}$ and its evolution with time. The behavior of $dr/d\ln a$ deserves attention with regard to the coincidence problem. Clearly we see that the positive coupling obtained from the best fit data leads slower change of r which makes the coincidence problem less acute when compared with the case without interaction shown by the red dotted line in Fig 2a. When we choose DE EoS to be ω_{II} and constant, results are shown in Fig 2b,c. From the behaviors of the ratio $r = \rho_{DM}/\rho_{DE}$, for the constant DE EoS or ω_{II} , this phenomenological interaction cannot help to alleviate the coincidence problem. Besides, the negative coupling shown in Fig 1b for DE EoS ω_{II} leads negative ρ_{DE} in the early time. When DE EoS is constant ρ_{DE} is nearly zero even in the inflationary era, which makes it hard to count on this ρ_{DE} to explain the inflation. What's worse, the ρ_{DM} will be negative in the future. All these unphysical consequences are brought by the negative coupling from the fitting.

Coupling	EoS		Ω_b^0	ω_0	ω_1	Ω_{DM}^0	h_0	δ
$\delta H \rho_{DE}$	ω_I	Mean	$0.052^{+0.006}_{-0.006}$	$-0.82^{+0.30}_{-0.23}$	$-1.97^{+0.99}_{-1.64}$	$0.27^{+0.03}_{-0.03}$	$0.66^{+0.05}_{-0.04}$	$-0.28^{+0.13}_{-0.20}$
		Best-fitted	0.054	-1.18	1.21	0.24	0.63	0.03
	ω_{II}	Mean	$0.055^{+0.007}_{-0.006}$	$-1.00^{+0.45}_{-0.37}$	$-0.72^{+2.65}_{-3.68}$	$0.25^{+0.03}_{-0.03}$	$0.64^{+0.05}_{-0.05}$	$-0.22^{+0.12}_{-0.18}$
		Best-fitted	0.059	-1.26	2.03	0.23	0.62	-0.08
	constant ω		Ω_b^0	ω		Ω_{DM}^0	h_0	δ
		Mean	$0.055^{+0.007}_{-0.006}$	$-1.05^{+0.12}_{-0.13}$		$0.25^{+0.02}_{-0.02}$	$0.64^{+0.05}_{-0.04}$	$-0.19^{+0.10}_{-0.13}$
		Best-fitted	0.057	-1.00		0.24	0.63	-0.13

B. The coupling is proportional to the energy density of DM

Using the first parameterization of the DE EoS and choosing the coupling between DE and DM in proportional to the energy density of DM, we show the likelihood in Fig 3a and the behaviors of cosmological parameters in Fig 4a. Comparing lines of $dr/d\ln a$ for the interacting and noninteracting cases, it seems that near the present time the interaction between DE and DM makes the change of r a bit slower than the noninteracting case, which makes the coincidence problem less serious. However since the coupling is negative from the best fit, we encounter the negative ρ_{DE} in the early epoch of the universe. This sounds unphysical[36], although in some modified gravity theory negative value of ρ_{DE} was allowed[37]. The results for choosing DE EoS to be ω_{II} and constant are shown in Fig 4b and Fig 4c. It is clear from the behavior of r that the coincidence problem cannot be alleviated by choosing this phenomenological interaction form when DE EoS are of ω_{II} or constant. Moreover, negative ρ_{DE} at very early epoch appears again.




Figure 2: These figures show clearly the behaviors of different dark sectors during the late time evolution of universe with best-fitted data for coupling $\delta H \rho_{DE}$. The red dotted line denotes the noninteracting case. The left one corresponds to EoS ω_I , the middle one is EoS ω_{II} and the right one is constant EoS




Figure 3: The likelihood for coupling parameters when the interaction is chosen as $\delta H \rho_{DM}$.

Coupling	EoS		Ω_b^0	ω_0	ω_1	Ω_{DM}^0	h_0	δ
$\delta H \rho_{DM}$	ω_I	Mean	$0.047^{+0.013}_{-0.008}$	$-0.99^{+0.20}_{-0.17}$	$0.34^{+0.37}_{-0.92}$	$0.25^{+0.02}_{-0.02}$	$0.69^{+0.08}_{-0.09}$	$-0.020^{+0.010}_{-0.010}$
		Best-fitted	0.044	-1.02	0.60	0.26	0.71	-0.024
	ω_{II}	Mean	$0.048^{+0.012}_{-0.008}$	$-1.25^{+0.31}_{-0.31}$	$2.35^{+1.96}_{-2.05}$	$0.25^{+0.02}_{-0.02}$	$0.69^{+0.08}_{-0.09}$	$-0.019^{+0.010}_{-0.008}$
		Best-fitted	0.046	-1.28	2.65	0.24	0.69	-0.021
	constant ω		Ω_b^0	ω	Ω_{DM}^0	h_0	δ	
		Mean	$0.046^{+0.010}_{-0.008}$	$-0.90^{+0.08}_{-0.08}$	$0.25^{+0.02}_{-0.02}$	$0.71^{+0.07}_{-0.08}$	$-0.018^{+0.010}_{-0.008}$	
		Best-fitted	0.041	-0.90	0.25	0.73	-0.022	

C. The coupling is proportional to the total energy density of DE and DM

The results of the evolution of cosmological parameters for employing ω_I and expressing the interaction between DE and DM in proportional to the total energy density of DE and DM are shown in Fig 5a and Fig 6a. In Fig 6a we see that r changes a bit slower than that of the noninteracting case at very recent epoch, and this might help to alleviate the coincidence problem. However in the early epoch the negative coupling obtained from fitting makes ρ_{DE} negative again. Choosing DE EoS to be ω_{II} , besides the negative ρ_{DE} in the early time, ρ_{DM} will appear negative in the future. For constant DE EoS, ρ_{DE} can also be negative in the early epoch. Furthermore from Fig 5b,c, we see that the coincidence problem has not been alleviated.




Figure 4: These figures show clearly the behaviors of different dark sectors during the late time evolution of universe with best-fitted data for the coupling $\delta H \rho_{DM}$. The red dotted line denotes the noninteracting data. The left one corresponds to EoS ω_I , the middle one is EoS ω_{II} and the right one is constant EoS




Figure 5: The likelihood for coupling parameters when the interaction is chosen as $\delta H(\rho_{DM} + \rho_{DE})$.

Coupling	EoS		Ω_b^0	ω_0	ω_1	Ω_{DM}^0	h_0	δ
$\delta H(\rho_{DM} + \rho_{DE})$	ω_I	Mean	$0.046^{+0.012}_{-0.008}$	$-0.98^{+0.20}_{-0.17}$	$0.21^{+0.49}_{-0.87}$	$0.25^{+0.02}_{-0.02}$	$0.70^{+0.08}_{-0.09}$	$-0.018^{+0.009}_{-0.008}$
		Best-fitted	0.044	-1.07	0.71	0.25	0.71	-0.022
	ω_{II}	Mean	$0.048^{+0.013}_{-0.008}$	$-1.22^{+0.34}_{-0.34}$	$2.09^{+2.08}_{-2.24}$	$0.246^{+0.02}_{-0.02}$	$0.69^{+0.08}_{-0.09}$	$-0.017^{+0.010}_{-0.009}$
		Best-fitted	0.044	-1.25	2.39	0.25	0.71	-0.021
	constant ω		Ω_b^0	ω	Ω_{DM}^0	h_0	δ	
		Mean	$0.045^{+0.010}_{-0.008}$	$-0.91^{+0.08}_{-0.08}$	$0.25^{+0.02}_{-0.02}$	$0.71^{+0.08}_{-0.08}$	$-0.017^{+0.009}_{-0.008}$	
		Best-fitted	0.042	-0.91	0.25	0.72	-0.020	

D. The coupling is proportional to the linear combination of energy densities of DE and DM

If the coupling is in linear combination of energy densities of DE and DM, we can see easily from Fig 9, the coincidence problem gets even worse, since r changes even faster than that of the noninteracting case. Furthermore, we find the negative ρ_{DM} in the future for all the cases, which is unphysical.




Figure 6: These figures show clearly the behaviors of different dark sectors during the late time evolution of universe with best-fitted data for the coupling $\delta H(\rho_{DM} + \rho_{DE})$. The red dotted line denotes the noninteracting case. The left one corresponds to EoS ω_I , the middle one is EoS ω_{II} and the right one is constant EoS




Figure 7: The likelihood for coupling parameters when the interaction is $H(\delta_1 \rho_{DE} + \delta_2 \rho_{DM})$. The left one shows the likelihood for parameter δ_1 and the right one show the likelihood for parameter δ_2 . The EoS is chosen as ω_I for these figures.

Coupling	EoS		Ω_b^0	ω_0	ω_1	Ω_{DM}^0	h_0	δ_1	δ_2
$H(\delta_1 \rho_{DE} + \delta_2 \rho_{DM})$	ω_I	Mean	$0.043^{+0.007}_{-0.006}$	$10.97^{+8.66}_{-9.42}$	$-473.12^{+220.46}_{-611.05}$	$0.30^{+0.05}_{-0.05}$	$0.72^{+0.06}_{-0.05}$	$-13.33^{+4.57}_{-4.13}$	$0.39^{+0.05}_{-0.05}$
		Best-fitted	0.037	1.41	-42.10	0.27	0.77	-4.26	0.30
	ω_{II}	Mean	$0.048^{+0.012}_{-0.009}$	$1.55^{+1.64}_{-0.34}$	$-188.36^{+39.10}_{-82.24}$	$0.23^{+0.03}_{-0.03}$	$0.69^{+0.08}_{-0.09}$	$-13.67^{+10.10}_{-5.90}$	$0.56^{+0.09}_{-0.08}$
		Best-fitted	0.043	0.43	-34.65	0.24	0.71	-4.55	0.53
	constant ω	Mean	$0.032^{+0.002}_{-0.002}$	$-2.31^{+0.39}_{-0.61}$	$0.24^{+0.02}_{-0.02}$	$0.82^{+0.03}_{-0.03}$	$-5.56^{+1.33}_{-1.87}$	$0.61^{+0.22}_{-0.18}$	
		Best-fitted	0.032	-1.40	0.24	0.82	-2.47	0.49	

E. The coupling is proportional to the product of energy densities of DE and DM

In Fig 11, we have shown the results by choosing the interaction as the product of energy densities of DE and DM ω_I . Choosing DE EoS ω_I , it is clear that with this kind of interaction between DE and DM, there is a slower change of r as compared to the noninteracting case. This means that the period when energy densities of DE and DM are comparable is longer compared to the noninteracting case. Thus it is not so strange that we now live in the coincidence state of the universe. In this sense the coincidence problem is less acute when compared with the case without interaction. Adopting this kind of interaction form, we have not met unphysical problems as mentioned




Figure 8: The likelihood for coupling parameters when the interaction is $H(\delta_1\rho_{DE} + \delta_2\rho_{DM})$. The left one shows the likelihood for parameter δ_1 and the right one show the likelihood for parameter δ_2 . The EoS is chosen as ω_{II} for these figures.




Figure 9: These figures show clearly the behaviors of different dark sectors during the late time evolution of universe with best-fitted data for the coupling $H(\delta_1\rho_{DE} + \delta_2\rho_{DM})$. The red dotted line denotes the noninteracting case. The left one corresponds to EoS ω_I , the middle one is EoS ω_{II} and the right one is constant EoS

above. Choosing DE EoS as ω_{II} , we see that in most period, r changes slower than that of the noninteracting case. Only very recently r changes a bit faster for the interacting case. This will not change the qualitative behavior of the period when energy densities of DE and DM are comparable, which is still longer compared to the noninteracting case. Thus the coincidence problem can become less serious. Choosing the interaction in the form of the product of densities of DE and DM, we have positive coupling from the fitting, which can help to avoid unphysical problems mentioned above. For constant DE EoS, we see that r changes a bit faster than that of the noninteracting case near the present epoch, which cannot help to alleviate the coincidence problem. However the positive coupling from fitting again avoids unphysical problems for the evolution of cosmological parameters.

Coupling	EoS		Ω_b^0	ω_0	ω_1	Ω_{DM}^0	h_0	δ
$\lambda\rho_{DE}\rho_{DM}$	ω_I	Mean	$0.036^{+0.003}_{-0.004}$	$-0.93^{+0.11}_{-0.11}$	$0.48^{+0.16}_{-0.25}$	$0.25^{+0.018}_{-0.018}$	$0.78^{+0.047}_{-0.037}$	$0.012^{+0.004}_{-0.004}$
		Best-fitted	0.036	-0.98	0.68	0.25	0.78	0.0064
	ω_{II}	Mean	$0.040^{+0.003}_{-0.003}$	$-1.53^{+0.35}_{-0.35}$	$4.90^{+2.13}_{-2.26}$	$0.24^{+0.020}_{-0.018}$	$0.74^{+0.028}_{-0.027}$	$0.018^{+0.020}_{-0.007}$
		Best-fitted	0.041	-1.52	4.93	0.23	0.73	0.016
	constant		Ω_b^0	ω		Ω_{DM}^0	h_0	δ
		Mean	$0.039^{+0.002}_{-0.003}$	$-0.84^{+0.08}_{-0.09}$	$0.25^{+0.02}_{-0.02}$	$0.74^{+0.03}_{-0.03}$	$0.0063^{+0.0074}_{-0.0063}$	
		Best-fitted	0.038	-0.85	0.25	0.75	0.0025	




Figure 10: The likelihood for coupling parameters when the interaction is chosen as $\lambda\rho_{DM}\rho_{DE}$. The left one shows the EoS ω_I and the right one is ω_{II}




Figure 11: These figures show clearly the behaviors of different dark sectors during the late time evolution of universe with best-fitted data for the interaction $\lambda\rho_{DM}\rho_{DE}$. The red dotted line denotes the noninteracting case. The left one corresponds to EoS ω_I , the middle one is EoS ω_{II} and the right one is constant EoS

IV. CONCLUSIONS AND DISCUSSIONS

In this paper we have examined the effects of all possible phenomenological interactions between DE and DM on cosmological parameters and their efficiency on solving the coincidence problem. We have worked with two simple parameterizations of the DE EoS and the constant DE EoS. Using observational data coming from the new 182 Gold type Ia supernova samples, the shift parameter of the Cosmic Microwave Background given by the three-year Wilkinson Microwave Anisotropy Probe observations, and the baryon acoustic oscillation measurement from the Sloan Digital Sky Survey, we have performed a statistical joint analysis of different forms of phenomenological interaction between DE and DM. We have found that for the time-dependent DE EoS the model of the interaction in product of densities of DE and DM is the best model to alleviate the coincidence problem. For other interaction forms, the consequences of DE and DM interaction on cosmological parameters are very sensitive to the DE EoS. For the DE EoS in the form of ω_I , choosing the DE and DM interaction in proportional to the energy density of DE, we found the solution to solve the coincidence problem. We observed that other forms of the interaction either bring unphysical problems or cannot help to alleviate the coincidence problem. For the DE EoS ω_{II} , we found that except the interaction in terms of the product of densities of DE and DM, other phenomenological descriptions of interaction between DE and DM fail to alleviate the coincidence problem. For the constant DE EoS, none of the phenomenological models of interaction can help to alleviate the coincidence problem. It would be interesting to extend our analysis to more general parameterizations for both the interaction and the DE EoS available in the literature. Furthermore it is also of interest to include more observational data to constrain the interaction between DE and DM and study its consequences on the cosmological parameters.

Acknowledgments

This work was partially supported by the NNSF of China, Shanghai Education Commission, Shanghai Science and Technology Commission.

-
- [1] A. G. Riess *et al*, *Astron. J.* **116**, 1009 (1998); S. Perlmutter *et al*, *Astrophys. J.* **517**, 565 (1999); P. de Bernardis *et. al.*, *Nature* **404**, 955 (2000).
 - [2] S. Perlmutter *et al*, *Astrophys. J.* **598**, 102 (2003).
 - [3] D. J. Eisenstein, *et al.*, *Astrophys. J.* **633**, 560 (2005).
 - [4] D. N. Spergel *et al.*, *Astrophys. J. Suppl.* **170**, 377 (2007).
 - [5] A. G. Riess, *et al.*, astro-ph/0611572.
 - [6] T. Padmanabhan, *Phys. Rept.* **380**, 235 (2003); P. J. E. Peebles, B. Ratra, *Rev. Mod. Phys.* **75**, 559 (2003); V. Sahni, *Lect Notes Phys.* **653**:141 (2004) and references therein.
 - [7] L. Amendola, *Phys. Rev.* **D62**, 043511 (2000); L. Amendola and C. Quercellini, *Phys. Rev.* **D68**, 023514 (2003); L. Amendola, S. Tsujikawa and M. Sami, *Phys. Lett.* **B632**, 155 (2006).
 - [8] D. Pavon, W. Zimdahl, *Phys. Lett.* **B628**, 206 (2005).
 - [9] S. Campo, R. Herrera, G. Olivares and D. Pavon, *Phys. Rev.* **D74**, 023501 (2006); S. Campo, R. Herrera and D. Pavon, *Phys. Rev.* **D71**, 123529 (2005); G. Olivares, F. Atrio-Barandela and D. Pavon, *Phys. Rev.* **D71**, 063523 (2005).
 - [10] G. Olivares, F. Atrio-Barandela and D. Pavon, *Phys. Rev.* **D74**, 043521 (2006).
 - [11] B. Wang, Y. G. Gong and E. Abdalla, *Phys. Lett.* **B624**, 141 (2005).
 - [12] B. Wang, C.-Y. Lin and E. Abdalla, *Phys. Lett.* **B637**, 357 (2006).
 - [13] B. Wang, J. Zang, C.-Y. Lin, E. Abdalla and S. Micheletti, *Nucl. Phys.* **B778**, 69 (2007).
 - [14] G. Mangano, G. Miele and V. Pettorino, *Mod.Phys.Lett.A* **18**, 831(2003).
 - [15] S. Das, P. S. Corasaniti and J. Khoury, *Phys. Rev.* **D73**, 083509 (2006).
 - [16] R. Bean, E. E. Flanagan, M. Trodden, arXiv:0709.1124; R. Bean, E. E. Flanagan, M. Trodden, arXiv:0709.1128; M. Manera, D. F. Mota arXiv:astro-ph/0504519; N. J. Nunes, D. F. Mota arXiv:astro-ph/0409481
 - [17] W. Zimdahl, *Int. J. Mod. Phys.* **D14**, 2319 (2005).
 - [18] E. Abdalla and B. Wang, *Phys. Lett.* **B651**, 89 (2007).
 - [19] C. Feng, B. Wang, Y. G. Gong, R. K. Su, *JCAP* **09**, 005 (2007).
 - [20] Z. K. Guo, N. Ohta and S. Tsujikawa, *Phys. Rev.* **D76**, 023508 (2007).
 - [21] O. Bertolami, F. Gil Pedro and M. Le Delliou, *Phys. Lett.* **B654**, 165 (2007). O. Bertolami, F. Gil Pedro and M. Le Delliou, arXiv:0705.3118v1.
 - [22] M. Kesden and M. Kamionkowski, *Phys. Rev. Lett.* **97**, 131303 (2006); M. Kesden and M. Kamionkowski, *Phys. Rev.* **D74**, 083007 (2006).
 - [23] E. Abdalla, L. Raul W. Abramo, L. Sodre Jr., B. Wang, arXiv:0710.1198 [astro-ph].
 - [24] L. Amendola, D. Tocchini-Valentini, *Phys. Rev. D* **64** (2001) 043509; G. W. Anderson, S. M. Carroll, astro-ph/9711288.
 - [25] S. Das, P. S. Corasaniti, J. Khoury, *Phys. Rev. D* **73** (2006) 083509.
 - [26] W. Zimdahl, D. Pavj on, L. P. Chimento, *Phys. Lett. B* **521** (2001) 133; L. P. Chimento, A. S. Jakubi, D. Pavj on, W. Zimdahl, *Phys. Rev. D* **67** (2003) 083513; S. del Campo, R. Herrera, D. Pavj on, *Phys. Rev. D* **70** (2004) 043540; D. Pavj on, W. Zimdahl, *Phys. Lett. B* **628** (2005) 206.
 - [27] B. Wang, C.-Y. Lin, D. Pavon, E. Abdalla, *Phys. Lett.* **B662**, 1 (2008).
 - [28] D. Pavon, B. Wang, arXiv:0712.0565.
 - [29] U. Alam, V. Sahni, and A. Starobinsky, *JCAP* **0406**, 008 (2004); Y. G. Gong, *Class. Quant. Grav.* **22**, 2121 (2005); Y. Wang and M. Tegmark, *Phys. Rev. D* **71**, 103513 (2005); Y. Wang and P. Mukherjee, *Astrophys. J.* **606**, 654 (2004); R. Daly and S. Djorgovski, *Astrophys. J.* **612**, 652 (2004); U. Alam, V. Sahni, T. Saini, and A. Starobinsky, *Mon. Not. Roy. Astron. Soc.* **354**, 275 (2004); T. Choudhury and T. Padmanabhan, *Astron. Astrophys.* **429**, 807 (2005).
 - [30] D. N. Spergel, *et al.*, 2007, *ApJS*, **170**, 377.
 - [31] A. G. Riess, *et al.*, 2006, arXiv: astro-ph/0611572.
 - [32] D. J. Eisenstein, *et al.*, 2005, *ApJ*, **633**, 560.
 - [33] O. Elgaroy, T. Multamaki, 2007, arXiv:astro-ph/0702343.
 - [34] Y. Wang, P. Mukherjee, 2007, arXiv:astro-ph/0703780.
 - [35] W. L. Freedman, *et al.*, 2001, *ApJ*, **553**, 47.
 - [36] L. Amendola, G. Campos, R. Rosenfeld, astro-ph/0610806.
 - [37] L. Amendola, D. Polarski, S. Tsujikawa, astro-ph/0603703.
 - [38] M. Doran, C. M. Muller, astro-ph/0311311.